

PRODUCT DATA SHEET

Donor Horse Serum

Description

Donor horse serum (DHS) is obtained from closed donor horse herds that receive regular veterinary inspection and care. The donor animals are all gelded draft horses in order to limit lot-to-lot hormone level variations. They are fed a special diet and are fasted prior to bleeding to reduce lipid concentrations. DHS will support the growth of a wide variety of mammalian cells, providing a cost-effective alternative for fetal bovine serum. DHS is often used in bovine virus research applications since it does not contain bovine antibodies.

Collection and Processing

Atlanta Biologicals consistently supplies DHS lots with excellent cell growth characteristics, low endotoxin and low hemoglobin values.

Origin:

The DHS used in Atlanta Biologicals' manufacturing process meets all of the USDA requirements for animal products. All of our DHS is traceable back to the date and location of collection. In addition, all of Atlanta Biologicals' serum used in manufacturing must meet our strict quality requirements for raw material.

Raw Material Processing:

Animal blood is aseptically collected from prescreened donor animals that are maintained in controlled herds. The whole blood is allowed to clot at refrigerated temperatures. Serum is then carefully removed from the clot after centrifugation at refrigerated temperatures, to avoid contamination by red blood cells. This raw serum product is immediately placed into bottles and frozen for delivery to our manufacturing facility. The product remains frozen throughout the entire shipping and receiving process, from the raw processing site to our manufacturing facility. This rapid processing ensures that endotoxin levels in the serum remain low and that the growth promoting qualities of the serum remain at their peak levels.

Filtration:

Approved lots of raw serum are thawed under controlled conditions and sterile filtered by an in-line process that uses a series of filters in descending pore size, utilizing 0.2 μ m filters for the final filtration step. Filling takes place in a laminar flow hood certified to maintain Class 100 conditions. The filling room is maintained under positive pressure with HEPA-filtered air. The serum is aseptically dispensed into gamma irradiated, sterile PETG or PETE bottles. Filled containers are immediately labeled and frozen, and then maintained at temperatures less than -5°C to preserve the product quality.

This product is manufactured for research and development purposes only. It is not intended for any human or animal diagnostic, therapeutic or other clinical uses. It is also not for agricultural, food, drug, cosmetic or household use. The use of these products must be supervised by a person technically qualified to handle potentially hazardous material.

Quality Control Testing

Chemical Analyses:

The Osmolality (vapor pressure method) and pH are measured on instruments that are calibrated daily using reference solutions traceable to National Institute of Standards and Technology Reference Materials.

Hemoglobin content of the serum is measured spectrophotometrically.

Endotoxin content is measured using the Limulus ameobocyte lysate (LAL) gel-clotting assay.

Biochemical Profile:

Total Protein	Total Bilirubin	Blood Urea Nitrogen (BUN)	Sodium/Potassium Ratio
Albumin	Iron	Creatinine	Chloride
Globulin	UIBC	BUN/Creatinine Ratio	Calcium
A/G ratio	Cholesterol	Uric Acid	Phosphorus
IgG	Triglycerides	Sodium	Magnesium
ALT/SGPT	Glucose	Potassium	Bicarbonate
Alkaline Phosphatase			

Microbiological Testing:

Each lot of serum is tested to confirm the absence of bacterial or fungal contamination using modified methods referenced in the U.S. Pharmacopeia (USP).

Each lot of serum is tested to confirm the absence of mycoplasma contamination to the limit of detection with the methods used. The large-volume method of Barile and Kern is used to detect mycoplasma that can be cultivated in media. Three different media are inoculated with the serum sample and grown under both aerobic and anaerobic conditions. Non-cultivable mycoplasma are detected by passage of the sample on an indicator cell line and staining with a DNA-fluorochrome.

Storage and Handling

The DHS is supplied in gamma irradiated, sterile PETG or PETE bottles. We recommend that the serum be stored frozen at a temperature of -5°C to -20°C . Multiple freeze-thaw cycles of the serum should be avoided as this may lead to deterioration of the product. If intermittent usage of the product is anticipated, we recommend use of either our smaller package sizes or dividing the serum into smaller aliquots suitable for single use. Always use aseptic techniques when handling the serum and aliquot into sterile containers.

Shipping

Serum is shipped frozen by next day air in insulated containers packed with dry ice.

This product is manufactured for research and development purposes only. It is not intended for any human or animal diagnostic, therapeutic or other clinical uses. It is also not for agricultural, food, drug, cosmetic or household use. The use of these products must be supervised by a person technically qualified to handle potentially hazardous material.